


CHE 103e



OFFICE OF THE DEPUTY PRINCIPAL
ACADEMICS, STUDENT AFFAIRS AND RESEARCH

UNIVERSITY EXAMINATIONS

2018/2019 ACADEMIC YEAR

FIRST YEAR SECOND SEMESTER REGULAR EXAMINATION

**FOR THE DEGREE OF BACHELOR OF
EDUCATION SCIENCE**

COURSE CODE: CHE 103e

**COURSE TITLE: INTRODUCTION TO
THERMODYNAMICS AND
KINETICS**

DATE: 24TH APRIL, 2019

TIME: 2.00 PM – 5.00 PM

INSTRUCTION TO CANDIDATES

- SEE INSIDE



THIS PAPER CONSISTS OF 4 PRINTED PAGES

PLEASE TURN OVER

CHE 103e: INTRODUCTION TO THERMODYNAMICS AND KINETICS

STREAM: BED (Science)

DURATION: 3 Hours

INSTRUCTIONS TO CANDIDATES


- i. Answer ALL questions.
- ii. Use the following physical constants where applicable:

Physical Constants

$R = 0.08206 \text{ atm L K}^{-1} \text{ mol}^{-1}$ or $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$, $1^\circ\text{C} = 273\text{K}$, $\text{CO}_{2(g)}$, $a = 3.592 \text{ a}$
 $(\text{L}^2\text{-atm/mol}^2)$ and $b = 0.04267 \text{ (L/mol)}$, $N = 14$

Question One

- a) Define the following terms:
 - i. Ideality (1 Mark)
 - ii. Thermodynamics (1 Mark)
 - iii. Closed and isothermal system (1 Mark)
 - iv. Extensive Variable (1 Mark)
 - v. Isochoric process (1 Mark)
 - vi. Kinetic theory of matter (1 Mark)
 - vii. Collision Theory (1 Mark)
- b) What is an ideal gas? (1 Mark)
- c) State two basic properties of gases which differentiate gases from liquids and solids (2 Marks)
- d) State four main features of the ideal gas kinetic molecular theory. (2 Marks)
- e) Why do real gases deviate from ideal behaviour? Give reasons why (2 Marks)
- f) Consider a sample of 1.000 mol of $\text{CO}_{2(g)}$ confined to a volume of 3.000 L at 0°C . Calculate the pressure of the gas using:
 - (i) The ideal-gas equation, and (2 Marks)
 - (ii) The van der Waals equation. (2 Marks)
- g) Explain what is happening in the graph below (2 Marks)

**Question Two**

- Differentiate between an ideal and a real gas. (1 Mark)
- Fifty grams of N_2 occupies a volume of 750 mL at 298.15 K. Assuming the gas behaves ideally, calculate the pressure of the gas in atm. (3 Marks)
- Using equations differentiate between expansion work and free expansion. (2 Marks)
- Prove that the relationship between heat capacity at constant pressure (C_p) and heat capacity at constant volume (C_v) is given by $C_p - C_v = R$ (3 Marks)
- Calculate the heat required to increase the temperature of gaseous O_2 from $0^\circ C$ to $100^\circ C$.
 - At constant P (2 Marks)
 - At constant V (2 Marks)
 - Account for the difference in heat absorbed in (i) and (ii). (1 Mark)
- Calculate the minimum work done at $50^\circ C$ on 5 moles of CO_2 to form a precipitate from a volume of 50 L to a volume of 1 L when CO_2 is considered as a perfect gas. (3 Marks)
- Argon gas at 5 atm expands reverse adiabatically to twice ($5\times$) its initial volume. Calculate its final pressure given that $\gamma = 5 / 3$. (3 Marks)

Question Three

- 1 mole of methane at $200^\circ C$ and 10 atm expands adiabatically and reversibly until its temperature was $0^\circ C$. If methane is a perfect gas with heat capacity at constant pressure of $15 J mol^{-1} K^{-1}$, calculate:
 - Work done on methane (3 Marks)
 - Final pressure of methane gas. (2 Marks)

- b) From the thermodynamic definition of enthalpy, $H=U+PV$, prove that $\Delta H=q_p$ (3 Marks)
- c) Using suitable equations and examples, differentiate between
- (i) Unimolecular reactions (2 Marks)
 - (ii) Bimolecular reactions (2 Marks)
- d) State the laws of thermochemistry. (2 Marks)
- e) State Hess law. (1 Mark)

Question Four

- a) With relevant examples in chemical equations, differentiate between homogeneous and heterogeneous catalysis. (2 Marks)
- b) Define the following terms:
- (i) Limiting enthalpy of solution (1 Mark)
 - (ii) Standard reaction enthalpy (1 Mark)
- c) The reaction, $2\text{NO}_{(g)} \leftrightarrow \text{N}_{2(g)} + \text{O}_{2(g)}$, has a value of $K= 2400$ at 2000 K. If 0.61 g of NO are put in a previously empty 3.00 L vessel, calculate the equilibrium concentrations of NO, N_2 , and O_2 . (3 Marks)
- d) Derive the integrated rate equations for first (1st) order reactions. (2 Marks)
- e) Give two practical applications of catalysts in real life. (2 Marks)
- f) Calculate the standard enthalpy change for the reaction: (4 Marks)



The relevant enthalpy changes of formation are;

$$\Delta H_f^\theta[\text{NaHCO}_3(s)] = -950.8 \text{ kJmol}^{-1}$$

$$\Delta H_f^\theta [\text{Na}_2\text{CO}_3(s)] = -1130.7 \text{ kJmol}^{-1}$$

$$\Delta H_f^\theta[\text{CO}_2(g)] = -393.5 \text{ kJmol}^{-1}$$

$$\Delta H_f^\theta[\text{H}_2\text{O}(l)] = -285.8 \text{ kJmol}^{-1}$$
